



Insect  
Monitoring  
Systems

Trece, Inc.  
PO Box 129  
Adair, OK 74330

## PHEROCON® NAVAL ORANGEWORM (NOW) TRAP - INSTRUCTIONS FOR USE

- Trap Location:** In full shade, 1 to 3 feet inside dripline of tree. Do not “bury” trap in center of tree canopy.
- Trap Density:** Never use less than 2 traps. Follow these guidelines according to orchard size:
- 20 acres or less: 1 trap per 2 1/2 to 5 acres
  - 21 to 40 acres: 1 trap per 10 acres
  - 41 to 80 acres: 1 trap per 15 to 20 acres
  - More than 80 acres: 1 per per 20 acres.
- Reading Frequency:** Count eggs twice weekly. Continue to monitor for signs of egg hatching.
- Trap Servicing:** Inspect bait for freshness when reading traps.
- Bait Replacement:** Replace bait every 6 weeks, or more often if it is degraded by excess moisture or other factors that affect freshness.

Distributed by:  
Great Lakes IPM, Inc.  
10220 E Church Rd.  
Vestaburg, MI 48891

800-235-0285  
989-268-5693  
FAX: 989-268-5311

[www.greatlakesipm.com](http://www.greatlakesipm.com)  
Email:  
[glipm@greatlakesipm.com](mailto:glipm@greatlakesipm.com)

