

BIOCONTROL



ISOMATE*-P PHEROMONE

ACTIVE INGREDIENTS:

3, 13 Octadecadien-1-yl Acetate	90.0%
INERT INGREDIENTS	10.0%
TOTAL	100.0%

Keep out of reach of children

WARNING

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

STATEMENT OF PRACTICAL TREATMENT

Avoid inhalation of vapors or contact with liquid ingredients. Harmful if swallowed. Avoid contact with eyes. In case of eye contact, immediately flush with water. Get medical treatment if irritation persists. Wash hands thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not discharge into lakes, streams, ponds, or public waters unless in accordance with NPDES permit. For guidance, contact your regional office of the Environmental Protection Agency.

PACIFIC BIOCONTROL CORPORATION

14615 NE 13th Court, Suite A, Vancouver, WA 98685 U.S.A.
Telephone (360) 571-2247 or 1-800-999-8805

MADE IN JAPAN

EPA Est. No: 47265-JP-01
EPA Reg. No: 53575-17

NET CONTENTS: 500 Dispensers Units

One dispenser contains 0.0014 fl oz or 37.8 mg
Total content of package: 0.7 fl oz or 18.9 gm

ISOMATE® is a registered Trademark of Pacific Biocontrol Corporation

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

Pesticide Storage	Store unopened package at temperatures below 40°F in a dry location. Product may be stored in cold storage facilities used for food storage.
Pesticide Disposal	Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State, and Local procedures.
Container Disposal	Dispose of empty dispensers by burning or burying with prunings in winter. If burned, stay out of smoke. Dispose of dispensers according to applicable Federal, State, and Local procedures. Foil envelopes can be disposed of as household refuse.
General	Consult Federal, State, or local disposal authorities for approved alternative procedures.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

It is **critical** that ISOMATE-P be applied as directed.

- 1. Crop** Peach, almond, nectarine, cherry, prune, plum, apricot.
- 2. Pest** Peach Tree Borer (*Synanthedon exitiosa*).
- 3. Rate** Minimum of 100 dispensers per acre or 250 dispensers per hectare. Attach one dispenser per tree. Maximum of 250 dispensers per acre or 625 dispensers per hectare.
- 4. Application** Attach the dispensers onto branches above the mid point of the tree.
- 5. Timing** Apply prior to moth emergence in spring.
- 6. Precautions** Isomate-P prevents Peach Tree Borer from mating. However, if a major source of mated female moths of this species is present adjacent to the treated field, migration of these moths may significantly reduce the level of control achieved. Sources are likely to be untreated host trees within 300 yards of the treated field. This can be overcome by:
 - a. Treatment of **entire blocks** and not just sections of large conventionally treated orchards which frequently serve as sources of mated females.
 - b. Treatment of infestation sources with Isomate-P (e.g., a strip at least 50 yards wide nearest the treated field).
 - c. Treatment of infestation source with an effective insecticide. Other pests must be monitored and controlled by timely intervention with conventional insecticides.

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. The Manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.